## WE CLAIM:

An enteral composition comprising:

a protein source consisting of hydrolyzed whey protein;

a carbohydrate source; and

a lipid source including a mixture of medium and long chain triglycerides, the enteral composition having a caloric density of at least 1.4 kcal/mL, wherein the composition provides a ratio of non-protein calories per gram nitrogen of at least 90:1.

- 2. The enteral composition of Claim 1 wherein the lipid source comprises approximately 20% to 50% of the calorie distribution of the composition. 10
  - The enteral composition of Claim 1 including 100% of U.S. RDA of 3. vitamins and minerals in approximately 1500 kcal.
- The enteral composition of Claim 1 wherein the protein source 4. 15 comprises approximately 15% to about 20% of the calorie distribution of the composition.
  - 5. The enteral composition of Claim 1 wherein the composition includes per 1500 kcal of composition:

a zinc source providing from approximately 28.5 to 43.5 mg; a vitamin C source providing from approximately 405 to 615 mg; a selenium source providing from approximately 60 to 90 mg; a taurine source providing from approximately 120 to 180 mg; and a L-carnitine source providing from approximately 120 to 180 mg.

The enteral composition of Claim 1 further including a source of beta-6.

-19-

5

20

25

carotene.

Int or

5

7. A method for providing nutrition to a metabolically stressed patient comprising the step of administering to the patient a therapeutically effective amount of a composition comprising:

a protein source comprising approximately 15% to about 20% of the calorie distribution of the composition, the protein source consisting of hydrolyzed protein;

a carbohydrate source;

a lipid source;

the enteral composition having a caloric density of at least 1.4 kcal/mL; and the composition provides a ratio of non-protein calories per gram nitrogen of at least approximately 90:1.

- 8. The method of Claim 7 wherein the lipid source comprises approximately 20% to 50% of the calorie distribution of the composition.
  - 9. The method of Claim 7 wherein the composition includes 100% of U.S. RDA of vitamins and minerals in approximately 1500 kcal.
- 20 The method of Claim 7 wherein the composition is fed through a tube to the patient.
  - 11. The method of Claim 7 wherein the composition contains approximately 0.37% of the calories as cysteine.
  - 12. The method of Claim 7 wherein the composition includes per 1500 kcal of composition:

25

15

5

a zinc source providing from approximately 28.5 to 43.5 mg; a vitamin C source providing from approximately 405 to 615 mg; a selenium source providing from approximately 60 to 90 mg; a taurine source providing from approximately 120 to 180 mg; and a L-carnitine source providing from approximately 120 to 180 mg.

13. The method of Claim 7 wherein the composition further includes a source of beta- carotene.

14. An enteral composition for a metabolically stressed patient comprising: about 15% to about 20% of the calorie distribution of the composition consisting essentially of hydrolyzed whey protein;

a carbohydrate source comprising at least 35% of the composition; a lipid source comprising at least 20 by weight of the composition; and the composition having a caloric density of at least 1.4 kcal/mL and a ratio of non-protein calories per gram of nitrogen of at least about 90:1.

- 15. The enteral composition of Claim 14 which includes, per 1500 kcal: a zinc source providing from about 28.5 to about 43.5 mg zinc;

  20 a vitamin C source providing about 405 to 615 mg vitamin C;

  a selenium source providing about 60 to about 90 mg selenium;

  a taurine source providing about 120 to about 180 mg taurine; and

  a L-carnitine source providing about 120 to about 180 mg L-carnitine.
- 25 16. The enteral composition of claim 14 which has a caloric density of about 1.4 to about 1.8 kcal/mL.
  - 17. The enteral composition of Claim 14 wherein the composition further

comprises from about 0.1% to 2.0% free amino acids.

18. The enteral composition of Claim 14 including at least 0.1% free amino acid.

5

10

15

19 A method for providing nutrition to a metabolically stressed patient comprising the step of administering to the patient a therapeutically effective amount of a composition comprising:

a protein source comprising approximately 15% to about 20% of the calorie distribution of the composition, the protein source consisting essentially of hydrolyzed whey protein;

a carbohydrate source;

a lipid source;

the enteral composition having a caloric density of at least 1.4 kcal/mL; and the composition provides a ratio of non-protein calories per gram nitrogen of at least approximately 90:1.

20. The method of Claim 19 wherein the composition includes 100% of U.S.RDA of vitamins and minerals in approximately 1500 kcal.

20

25

21. The method of Claim 19 wherein the composition includes per 1500 kcal of composition:

a zinc source providing from approximately 28.5 to 43.5 mg;

a vitamin C source providing from approximately 405 to 615 mg;

a selenium source providing from approximately 60 to 90 mg;

a taurine source providing from approximately 120 to 180 mg;

a L-carnitine source providing from approximately 120 to 180 mg; and

5

a source of beta- carotene.

22. The method of Claim 19 wherein the composition further comprises at least 0.1% free amino acids.

-23-